



FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF CALIFORNIA

DIRECTIONS FOR USE ON BASIL, MARJORAM, ROSEMARY, AND SAVORY

DEVRINOL® 50-DF SELECTIVE HERBICIDE
EPA REG NO. 70506-36 **EPA SLN NO. CA-060003**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. This label must be in the possession of the user at the time of pesticide application. Read the label affixed to the containers for DEVRINOL 50-DF Selective Herbicide before applying. Follow all applicable directions, restrictions and precautions on the EPA registered label. The use of DEVRINOL 50-DF Selective Herbicide according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the containers for DEVRINOL 50-DF Selective Herbicide.

DIRECTIONS FOR USE

Basil, Marjoram, Rosemary and Savory (Summer and Winter): For control of Annual Grasses and Broadleaf Weeds.

Apply 2-4 lbs. of DEVRINOL 50-DF Selective Herbicide per broadcast acre as a preplant/preemergence application. Apply with ground spray equipment only, in 20-100 gallons of water per acre. Apply to a weed free surface. Shallow incorporate no deeper than seeding depth or sprinkler irrigate within 24 hours using sufficient water to wet the soil to a depth of 2 to 4 inches. Use the lower rate on light soil (coarse textured- sandy) and the higher rate on heavy soil (fine textured-clay). A list of weeds controlled is provided on the second page of this label.

DEVRINOL 50-DF Selective Herbicide is not recommended for use on soil containing more than 10% organic matter. Spraying on ground heavily covered with leaves or trash may adversely affect weed control. Do not apply to frozen ground.

Do not apply more than once per season.

Do not apply this product through any type of irrigation system.

There is no restriction between last application and harvest.

CROP ROTATION RESTRICTIONS

- Do not rotate to leafy vegetables within 2 months (60 days) following a DEVRINOL 50-DF Selective Herbicide treatment at rates not exceeding 2 lbs ai/A.
- Do not rotate to cereal grains within 6 months (180 days) following a DEVRINOL 50-DF Selective Herbicide treatment at rates not exceeding 2 lbs ai/A.

**DEVIRINOL 50-DF SELECTIVE HERBICIDE - EPA REG NO. 70506-36
EPA SLN NO. CA-060003**

- Do not rotate to any other crop not specified on the product(s) label within 12 months (365 days) following a DEVIRINOL 50-DF Selective Herbicide treatment at rates not exceeding 2 lbs ai/A.

WEEDS CONTROLLED BY 2-4 LB DEVIRINOL 50-DF Selective Herbicide (1-2 LB AI/ACRE)

NOTE: DEVIRINOL 50-DF Selective Herbicide does not control established weeds.

Barnyardgrass	Junglerice
Carpetweed	Lambsquarters
Common Purslane	Panicum
Crabgrass	Purple Cudweed
Fall Panicum	Redroot Pigweed
Florida Pusley	Smooth Pigweed
Giant Foxtail	Stinkgrass
Goosegrass	Witchgrass
Green Foxtail	Yellow Foxtail
Johnsongrass (seedling)	

WEEDS CONTROLLED BY 4 LB DEVIRINOL 50-DF Selective Herbicide (2 LB AI/ACRE)

NOTE: DEVIRINOL 50-DF Selective Herbicide does not control established weeds.

Galinsoga

24(c) Registrant:



United Phosphorus, Inc.
630 Freedom Business Center
King of Prussia, PA 19406
1-800-438-6071 • www.upi-usa.com

CA approval 4/19/06
Rev. 4/14/11