



Recommendation

FIFRA SECTION 2(ee) RECOMMENDATION

CUPROFIX[®] ULTRA 40 DISPERSS[®]

EPA Registration No. 70506-201

Instructions for lower minimum application rates on various crops

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This recommendation is made as permitted under FIFRA Section 2(ee).

Observe all applicable directions, precautions, restrictions, and limitations on the EPA-approved label. This labeling contains supplemental instructions for use of this product that may not appear on the package label. This supplemental label and the federally registered label must be in the possession of the user at the time of pesticide application.

DIRECTIONS FOR USE:

CROP	DISEASE	<u>RATE</u> Lbs. Product/A	USE INSTRUCTIONS
Tomatoes	Anthracnose, Bacterial Speck, Bacterial Spot, Early Blight, Gray Leaf Mold, Late Blight, Septoria Leaf Spot	0.75 – 3	Begin applications when disease first threatens and repeat at 5 to 10 day intervals or as needed depending on disease severity. Use higher rates when conditions favor disease development.
Peppers	Anthracnose, Bacterial Spot, Cercospora Leaf Spot	0.75 – 3	Begin applications when conditions first favor disease development and repeat at 5 to 10 day intervals or as needed depending on disease severity. Use higher rates when conditions are favorable for disease development. Maneb 75DF can be tank mixed with Cuprofix Ultra 40 Disperss for added disease control.



Recommendation

Cucurbits	Alternaria Leaf Spot, Angular Leaf Spot, Anthracnose, Downy Mildew, Gummy Stem Blight, Powdery Mildew, Watermelon Bacterial Fruit Blotch (suppression)	1 – 2	Begin application when conditions are favorable for disease development. Repeat at 5 to 7 day intervals. Use shorter intervals when conditions are favorable for disease development. NOTE: Crop injury may result from application at shorter intervals. Discontinue use if injury occurs.
Citrus	Algal Spot, Melanose, Scab	1.75 – 8	Apply as pre-bloom and post-bloom sprays. Use higher rates when conditions favor disease development.
	Greasy Spot, Pink Pitting	0.75 – 5	Apply in summer on expanded new flush. Repeat on subsequent flushes where disease pressure is severe. Use higher rates when conditions favor disease development.
	Alternaria Brown Spot (suppression)	1.75 – 6.5	On susceptible varieties, apply when the first spring flush appears and each flush thereafter. Application to the fruit should start after two thirds of the petals have fallen and be repeated on a 21-day schedule. Note: When using the lower rates, use shorter spray intervals (7 to 14 days).
	Phytophthora Brown Rot, Septoria Spot	1.75 – 6	Begin application in fall before or just after the first rain and continue as needed. Apply to entire tree for Septoria, or just the lower 4 to 5 feet of the tree for Brown Rot. Also apply to bare ground 1 foot beyond skirt. Use higher rates when conditions favor disease development.



Recommendation

Citrus cont	Phytophthora Foot Rot	0.5 – 1	Mix with 1 gallon of water and paint trunks of trees from the soil surface to the lowest scaffold limbs. Apply in May prior to summer rains and/or in the fall prior to wrapping trees for freeze protection. Treatment serves as protection for up to 1 year, but does not cure existing infections. Note: Areas where microjet or low volume irrigation hit the tree trunk may require retreatment due to wash off.
	Citrus Canker (suppression only)	1.25 – 8	Spray flushes 7 to 14 days after shoots begin to grow. Young fruit may require an additional application. Number and timing of applications will be dependent upon disease pressure. Under heavy pressure, each flush of new growth should be sprayed.