



Special Local Need

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF OREGON

Ziram 76DF Fungicide

EPA Reg. No. 70506-173

EPA SLN No. OR-180001

This label is valid until December 31, 2023 or until otherwise amended, withdrawn, cancelled, or suspended.

ACTIVE INGREDIENT:

Ziram (Zinc dimethyldithiocarbamate) 76.0%

OTHER INGREDIENTS: 24.0%

TOTAL **100.0%**

**KEEP OUT OF REACH OF CHILDREN
PELIGRO DANGER**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

ENVIRONMENTAL HAZARDS
This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from target area. Do not contaminate water when disposing of equipment washwater or rinsate.

AGRICULTURAL USE REQUIREMENTS
Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, including plants, soil, or water, is:
Coveralls
Chemical resistant gloves made of any waterproof material
Shoes plus socks
Protective eyewear

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.



Special Local Need

RESTRICTIONS

Do not apply this product through any type of irrigation system.

Do not graze or feed cover crops from treated orchards.

FOR USE IN CONTROL OF EASTERN FILBERT BLIGHT

Crop	Pounds Per Acre	Disease(s)	Application Instructions
Filberts	3 – 8 lbs/A	Eastern Filbert Blight (EFB)	<p>Apply at bud break and continue at 2 week intervals until early May. Thorough coverage of all branches is essential. Use the higher rate if disease pressure is high and/or if crop canopy is dense. Do not apply more than 5 applications per crop cycle. Do not apply more than 30 lbs (22.8 lbs a.i.) per crop cycle.</p> <p>For ground applications, apply in a minimum of 20 gallons per acre. For aerial applications, apply in a minimum of 10 gallons of spray per acre. For both ground and aerial applications, always use sufficient water to provide thorough coverage.</p>

Rev. 01/31/2018