



Supplemental Label

Satellite[®] HydroCap herbicide (EPA Reg. No. 70506-230)

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Follow all applicable directions, restrictions and precautions on the federally-registered label.

This label and the entire **Satellite HydroCap herbicide** label, EPA Reg. No. 70506-230, must be in the possession of the user at the time of pesticide application.

This Supplemental Label expires March 1, 2020 and must not be used or distributed after this date.

CAUTION

KEEP OUT OF REACH OF CHILDREN

For Use in Forage Grasses (Cool-Season)

Application Method: Apply by air, ground, chemigation, or on dry bulk fertilizer.

Use Methods, Timings and Use Rates

For use in established Perennial Cool-Season Grasses including bentgrass, bluegrass (Kentucky), Bromegrass, fescue (fine, tall), orchardgrass, perennial ryegrass, timothy, and wheatgrass grown for forage, green chop, silage, hay production, and/or grown in pastures, rangeland, or Federal Conservation Reserve Program (CRP) land for livestock grazing.

Apply **Satellite HydroCap herbicide** to these and similar established (having 6 or more tillers per plant) grasses. Make application before target weeds germinate in the fall after the last cutting/mowing/grazing; in winter; in the spring; or in-season between cuttings.

Make a uniform broadcast application at 1.1 to 4.2 quarts per acre in a single application or sequentially at least 30 days apart. Do not apply more than 4.2 quarts of **Satellite HydroCap herbicide** per acre per year. Use the higher rate when weed pressure is high or when a longer duration of residual weed control is required.

Satellite HydroCap herbicide may cause temporary injury to cool-season forage grass stands. If application is made in the following situations the risk of crop injury may be increased: in periods of cold temperatures that limit normal crop growth; in extended cold temperatures that initiate winter dormancy; disease, extremely cold weather, drought, extensive frost heaving, salinity, low pH, or high pH may also weaken stands and make the crop more susceptible to herbicidal injury.

Mixed Stands of Established Cool-Season Forage Grasses and Alfalfa

Apply **Satellite HydroCap herbicide** to mixed stands of established alfalfa (defined as alfalfa planted in the fall or spring which has gone through a first cutting/mowing) and cool-season forage grasses. Apply in the fall after the last cutting/mowing/grazing, in winter, in the spring before weed germination, or in-season between cuttings.

Chemigation Application

Satellite HydroCap herbicide may be applied through sprinkler irrigation systems. Follow all directions, special instructions, and precautions in the Spraying Instructions section of the Satellite HydroCap Herbicide label.

Tank Mixes

Satellite HydroCap herbicide may be applied in a sequential use program with other herbicides labeled for this use, or applied as a tank mix with other registered herbicides that control emerged weeds. Applications of postemergence herbicides may cause crop injury, so proposed tank mixes must be tested on a small portion of the target crop to determine if damage is likely. Always perform a mixing test to check compatibility of **Satellite HydroCap herbicide** with potential tank mix partners and fertilizers. Follow all precautions and restrictions on the labels of all products applied in combination with **Satellite HydroCap herbicide**, and always follow the most restrictive label.

Physical incompatibility, reduced weed control, or crop injury may result from mixing Satellite HydroCap herbicide with other pesticides, additives, or fertilizers.

Preharvest Interval (PHI): there is no PHI for **Satellite HydroCap herbicide**-treated grass forage, green chop, silage, hay, pasture, or rangeland.

Restrictions

- Do not apply more than 4.2 quarts of **Satellite HydroCap herbicide** per acre per year.
- Do not graze or harvest mixed stand alfalfa/cool season forage grasses for forage and hay until 14 or more days after application.
- Do not apply **Satellite HydroCap herbicide** to mixed stands of cool-season forage grasses with other forage legumes besides alfalfa.
- Do not apply if surface water is present in the field.

Rev. 1/25/17