



Special Local Need

RESTRICTED USE PESTICIDE

RESTRICTED USE PESTICIDE due to carcinogenicity, potential for affecting fetal development, and high acute toxicity to humans. For retail sales to and use by Certified Applicators or persons directly under their supervision, and only for those uses covered by the Certified Applicators Certification.

Super Tin® 4L

Triphenyltin Hydroxide Flowable Fungicide

FOR USE ONLY IN THE STATE OF NORTH DAKOTA

This label is valid until December 31, 2022 or until otherwise amended, withdrawn, cancelled or suspended.

EPA Reg. No. 70506-212

EPA SLN No. ND-170007

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This FIFRA Section 24 (c) Special Local Need (SLN) label and the federal label must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label. Read the entire Directions for Use before using this product.

APPLICATION INSTRUCTIONS FOR CONTROL OF CERCOSPORA ON SUGAR BEET

USE RATE	APPLICATION INSTRUCTIONS
4-8 fl. oz./Acre	<p>Ground (Closed cabs only): Apply in at least 15 gallons of water. Full coverage of the foliage is necessary for the best results.</p> <p>Aerial (helicopter or fixed wing aircraft): Apply in 5-10 gallons of water.</p> <p>Ground or Aerial Applications: Diluted spray should be directed uniformly to all parts of the plant. Use lower gallonage when plants are small and increase volume with plant size. Use the lower specified rate for protective sprays and the higher specified rates later in the season or during high infection periods. Application should begin when leaf spot conditions appear or when the disease is in the area and repeated at 10 and 14 day intervals.</p> <p>Do not exceed 24 fl. oz./Acre per season.</p> <p>Do not treat within 7 days of harvest.</p> <p>Do not treat graze or feed beet tops to livestock.</p>