



Recommendation

FIFRA SECTION 2(ee) RECOMMENDATION FOR DISTRIBUTION AND USE IN ALABAMA, ARKANSAS, LOUISIANA, AND MISSISSIPPI

Froghorn™ Fungicide EPA Reg. No. 8033-127-70506

For suppression or control of Soil Borne diseases in Peanuts

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by US EPA.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This label and the entire Froghorn Fungicide container label, EPA Registration Number 8033-127-70506, must be in the possession of the user at the time of the application. Read the label affixed to the container for Froghorn Fungicide before applying. Carefully follow all precautionary statements and applicable use directions.

CROP	USE RATE PER ACRE	DISEASES	REMARKS
PEANUTS Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.	12.4 fl oz/A	SOIL BORNE: Cylindrocladium Black Rot (suppression) Rhizoctonia Limb Rot Rhizoctonia Pod Rot	For control of soil borne diseases, tankmix 12.4 fl oz/A of Froghorn and 4.0 fl oz/A of Tebuzol 3.6F fungicide. Under a 7-spray program, application timing of Froghorn for optimum control of Rhizoctonia Limb and Rod Rot should be applied during applications 3, 4, 5 and 6. Applications of chlorothalonil should be made prior to and following, (applications 1, 2, and 7) applications of Froghorn, to discourage development of resistant strains of fungi. For control of soil borne diseases within a leaf spot advisory schedule, apply Froghorn in the first advisory spray in July and continue applications at 14-day intervals. Froghorn must be carried by rainfall or irrigation into the root and pod zone for control of root and pod rots caused by <i>Sclerotium rolfsii</i> and <i>Rhizoctonia solani</i> . Drought conditions will decrease the effectiveness of Froghorn against the root and pod rots. Use Froghorn in conjunction with cultural practices that are known to reduce the severity of soil borne diseases, such as proper crop rotation practices.
<ul style="list-style-type: none"> Do not exceed a total application of 1.4 lbs a.i. thiophanate-methyl and 0.81 lbs a.i. tebuconazole per acre per year from all applications of all products containing these active ingredients. Do not feed hay or threshings, or allow livestock to graze in treated areas. Follow resistance management guidelines included on the Froghorn label. PHI: 14 days 			



Recommendation

Read the Limitation of Warranty and Liability on the container labeling before buying or using Froghorn Fungicide. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the container label.

Rev. 04/20/2018