



Recommendation

FIFRA SECTION 2(ee) RECOMMENDATION FOR DISTRIBUTION AND USE IN ALL STATES EXCEPT CALIFORNIA, NEW YORK AND TEXAS

STORM® HERBICIDE EPA Reg. No. 70506-59

FOR CONTROL OF PIGWEED IN PEANUTS AND SOYBEANS

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by USEPA

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read and follow all applicable instructions, restrictions, and precautions on the EPA-approved label. This label and the federally registered label must be in the possession of the user at the time of pesticide application.

Weeds Controlled in Peanuts and Soybeans (including glyphosate, triazine and ALS-resistant biotypes)	Scientific Name	1.5 pints per acre	
		Leaf Stage ^a (up to)	Maximum Height
Pigweed, Palmer	<i>Amaranthus palmeri</i>	6	<4"
, Redroot	<i>Amaranthus retroflexus</i>	6	2"
, Prostrate	<i>Amaranthus blitoides</i>	4	4"
, Smooth	<i>Amaranthus hybridus</i>	6	3"
, Spiny	<i>Amaranthus spinosus</i>	2	<2"

^a Do not count leaves as pairs; count each leaf separately. Do not count cotyledon leaves. Do not spray weeds in the cotyledon growth stage.

Consult the container label for instructions on tank mixes.

PEANUTS: Apply 1.5 pints of **Storm** herbicide per acre to peanuts preemergence at cracking stage (initiation of soil cracking, but before peanut emergence from the soil), or postemergence to peanuts to control susceptible weeds observing the labeled pre-harvest interval of 75 days.

SOYBEANS: Apply 1.5 pints of **Storm** herbicide per acre to soybeans to control susceptible weeds, observing the labeled pre-harvest interval of 50 days. To ensure optimum spray coverage of weeds, apply **Storm** herbicide to small actively growing weeds.