



Recommendation

FIFRA SECTION 2(ee) RECOMMENDATION FOR DISTRIBUTION AND USE IN THE STATES OF ALABAMA, ARKANSAS, GEORGIA, LOUISIANA, MISSISSIPPI, MISSOURI, NORTH CAROLINA, SOUTH CAROLINA, AND TENNESSEE

TRICOR® 4F HERBICIDE EPA Reg. No. 70506-68

FOR FALL APPLICATION TO CORN AND SOYBEANS

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by USEPA.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read and follow all applicable instructions, restrictions, and precautions on the EPA-approved label. This label and the federally registered label must be in the possession of the user at the time of pesticide application.

WEEDS CONTROLLED	RATE: PINTS TRICOR® 4F/A (Rate pounds ai/A)
Italian Ryegrass, Palmer Amaranth, Common Chickweed, Field Pennycress, Henbit, Marestalk, Winter Annual Mustard spp., Prickly Lettuce, Purple Deadnettle, Shepherd's Purse, Yellow Rocket, common dandelion seedlings	0.5 – 1.12 (0.25 – 0.56)
SPECIFIC DIRECTIONS	
<p>TriCor 4F Herbicide may be used alone or in combination with other registered herbicides as a fall applied ground broadcast application for burn-down and residual control. Apply TriCor 4F in the fall (September 1 – November 30) after harvest of the previous crop and before fall weeds emerge. Tillage may follow the application but do not exceed an incorporation in depth greater than 2-3 inches. This application will reduce weed cover before spring planting but is not intended to provide full weed control for the next cropping season. Additional residual and post-emergence herbicide applications will be required for an Integrated Weed Management Program in the following crop. The length of residual control will increase with the application rate of TriCor 4F. If emerged weeds are present and are greater than 2 inches in height or diameter, use an appropriate alternative post emergence herbicide in a tank mixture with TriCor 4F. To obtain maximum burn-down of existing weeds, use crop oil concentrate (COC) or an adjuvant in the tank mixture. Control of established common dandelion requires a tank mixture containing at least 1 pint/A of a 4 pound/gallon 2,4-D herbicide. Soybeans can be planted at any normal time following spring. Corn can also be planted at any normal time the following spring after fall TriCor 4F herbicide rates of 0.53 pints per acre or less. Corn can be planted at 4 or more months after fall TriCor 4F application at the rates on this document greater than 0.53 pints per acre.</p>	

Rev. 9/3/15