



# Supplemental Label

## Penncozeb® 80WP Fungicide

EPA Reg. No. 70506-183

### Instructions for Use on Cucurbits, Ginseng, Large and Small Tropical Fruit

**DIRECTIONS FOR USE:**

It is a violation of federal law to use this product in a manner inconsistent with its label. This supplemental label and the federally registered label must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

Crops Included	Diseases Controlled	Rate per Application Penncozeb 80WP Lbs./Acre	Remarks	Restrictions
Cucurbit Crop Group:  Chayote Chinese waxgourd Citron melon Cucumber Gherkin Gourd, edible <i>Momordica</i> spp. Muskmelon Pumpkin Squash, summer Squash, winter Watermelon	Alternaria Leaf Spot Anthracnose Cercospora Leaf Spot Downy Mildew Gummy Stem Blight Scab	2.0 - 3.0	Start applications when the plants are in the two-leaf stage and repeat at 7- to 10-day intervals. Use sufficient water and direct spray to provide thorough coverage of both upper and lower leaf surfaces. For aerial applications, the minimum spray volume is 2 gallons per acre.  Some cantaloupe varieties (i.e. Harvest Queen, Gold Star, Super Star, Sweet and Early, and Saticoy) are sensitive to Penncozeb 80WP fungicide. Consult State Cooperative Extension Service Specialist prior to use.	Do not apply more than 25.6 lbs. product (19.2 lbs. active ingredient) per acre per year.  Do not apply more than 8 applications per year.  Do not apply within 5 days of harvest.
Ginseng	Alternaria Blight	2.0	Start applications when disease first threatens and repeat every 7-10 days as needed. In Wisconsin, apply with ground equipment and a	Do not apply more than 24 lbs. product (18 lbs. active ingredient) per acre per year.  Do not apply more



# Supplemental Label

Crops Included	Diseases Controlled	Rate per Application Penncozeb 80WP Lbs./Acre	Remarks	Restrictions
			minimum of 80 gallons of water per acre.	<p>than 12 applications per year.</p> <p>Do not apply within 30 days of harvest.</p>
<p>Large Tropical Fruit:</p> <p>Canistel Mamey Sapote Mango Sapodilla Star Apple (caimito) White Sapote</p>	<p>Anthracnose Black Spot (Cercospora) Phytophthora Fruit Rot</p>	2.0 - 2.5	<p>Start applications at flowering and continue at 14- to 21- day intervals. Direct spray to crown and blossom area. Use 20 to 100 gallons water per acre.</p>	<p>Do not apply more than 37.3 lbs. product (28 lbs. active ingredient) per acre per year.</p> <p>Do not apply more than 14 applications per year.</p> <p>Applications may be made up to the day of harvest.</p>
<p>Small Tropical Fruit:</p> <p>Atemoya Cherimoya Custard apple Sugar apple Sweetsop</p>	Anthracnose	2.0 – 2.5	<p>Begin applications at flowering and continue at a 7-day retreatment interval.</p> <p>Applications made with aerial equipment must be made in a minimum spray volume of 10 gallons per acre.</p>	<p>Do not apply more than 35 lbs. product active (26.25 lbs. active ingredient) per acre per year.</p> <p>Do not apply more than 14 applications per year.</p> <p>Applications may be made up to the day of harvest.</p>

Consult your local Extension Service, Crop Advisor, or United Phosphorus, Inc. representative for additional information.

EPA Approved 9/30/10  
 NY Approved 5/6/11  
 CA Approved 4/29/11

Penncozeb is registered trademark of United Phosphorus, Inc.