



Special Local Need

FOR DISTRIBUTION AND USE ONLY WITHIN THE
STATE OF MICHIGAN

This label is valid until December 31, 2022 or until otherwise amended, withdrawn, cancelled, or suspended

TOPSIN[®] M WSB

THIOPHANATE-METHYL FUNGICIDE

EPA Registration No. 8033-125-70506

EPA SLN No. MI-180002

DIRECTIONS FOR USE IN THE PREVENTION/CONTROL OF WHITE MOLD and CYLINDROCARPON ROOT ROT ON GINSENG

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Disease	Lbs product per acre	Remarks
White Mold, Rhizoctonia Root And Crown Rot	1.0 (0.7 lb a.i.)	Apply as a foliar spray at 14-day intervals as needed.
Cylindrocarpon Root Rot	1.0 (0.7 lb a.i.)	Apply as a soil drench in 100 gallons water/A until runoff. Apply at 14-day intervals as needed.

Restrictions:

- Do not apply within 365 days of harvest.
- Do not apply more than 4 lbs product (2.8 lbs a.i.) per acre per year.
- Do not enter or allow worker entry into treated areas during the Restricted Entry Interval (REI) of 24 hours unless appropriate personal protection equipment (PPE) for early entry is worn.

This product is toxic to fish. Take the necessary precautions to prevent the product from entering surface water. Observe all restrictions, precautions, and limitations on the EPA-approved label. This label and the federally registered label must be in the possession of the user at the time of pesticide application.

Topsin M WSB may be applied by ground application only. Applications through any type of irrigation (chemigation) system are prohibited. To avoid the development of tolerant or resistant strains of fungi, different chemistry should be alternated with Topsin M WSB.

If more than one benzimidazole product is used, the total poundage of all such products must not exceed 3.5 lbs/A/season.

Rev. 04/30/2018

SOLD BY:
United Phosphorus, Inc.
630 Freedom Business Center, Suite 402
King of Prussia, PA 19406
www.upi-usa.com
1-800-438-6071