



Special Local Need

DEVIRINOL® 50-DF SELECTIVE HERBICIDE

DIRECTIONS FOR USE ON EGGPLANT GROWN WITH PLASTIC MULCH

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF GEORGIA

EPA Reg. No. 70506-36

EPA SLN NO. GA-120002

KEEP OUT OF REACH OF CHILDREN CAUTION

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. This label must be in the possession of the user at the time of pesticide application. Read the label affixed to the container for Devrinol 50-DF before applying. Follow all applicable directions, restrictions and precautions on the EPA registered label. The use of Devrinol 50-DF according to this supplemental labeling is subject to all use precautions and limitations imposed by the labels affixed to the containers for Devrinol 50-DF.

DIRECTIONS FOR USE For control of Annual Grasses and Broadleaf Weeds

CROP	RATE/ACRE OF DEVIRINOL 50DF	USE DIRECTIONS AND RESTRICTIONS
Eggplant grown with plastic mulch on plant beds – preplant	2-4 pounds (1-2 lb ai)	Apply to a weed free soil before the laying of plastic mulch. <u>Soil incorporated:</u> Soil should be well worked yet moist enough to permit a thorough incorporation to a depth of 2 inches. Incorporate the same day as applied before laying of plastic mulch using equipment which will result in uniform incorporation of the herbicide to desired depth. <u>Surface application:</u> If weed pressure is from small seeded annuals, apply to the surface of the bed immediately in front of the laying of the plastic mulch. Apply the plastic mulch over the treated soil the same day. Care must be taken to not move the soil on the treated bed when laying mulch. Do not exceed a maximum application rate of 4 lbs. per acre per crop cycle.

Rev. 4/5/12