



Special Local Need

**FOR DISTRIBUTION AND USE ONLY WITHIN THE
STATE OF TEXAS**

DIRECTIONS FOR USE ON ROSEMARY

DEVIRINOL[®] 50-DF SELECTIVE HERBICIDE
EPA REG NO. 70506-36 **EPA SLN NO. TX-100019**

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. This label must be in the possession of the user at the time of pesticide application. Read the label affixed to the containers for DEVIRINOL 50-DF before applying. Follow all applicable directions, restrictions and precautions on the EPA registered label. The use of DEVIRINOL 50-DF according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the containers for DEVIRINOL 50-DF.

For control of Annual Grasses and Broadleaf Weeds

Apply 2-4 lbs. of DEVIRINOL 50-DF Selective Herbicide per broadcast acre as a preplant/preemergence application. Apply with ground spray equipment only, in 20-100 gallons of water per acre. Apply to a weed free surface. Shallow incorporate no deeper than seeding depth or sprinkler irrigate within 24 hours using sufficient water to wet the soil to a depth of 2 to 4 inches. Use the lower rate on light soil (coarse textured- sandy) and the higher rate on heavy soil (fine textured-clay). A list of weeds controlled is provided on the second page of this label.

DEVIRINOL 50-DF Herbicide is not recommended for use on soil containing more than 10% organic matter. Spraying on ground heavily covered with leaves or trash may adversely affect weed control. Do not apply to frozen ground.

Do not apply more than once per season.

Do not apply this product through any type of irrigation system.

There is no restriction between last application and harvest.

CROP ROTATION RESTRICTIONS

- Do not rotate to leafy vegetables within 2 months (60 days) following a DEVIRINOL 50-DF treatment at rates not exceeding 2 lbs ai/A.
- Do not rotate to cereal grains within 6 months (180 days) following a DEVIRINOL 50-DF treatment at rates not exceeding 2 lbs ai/A.
- Do not rotate to any other crop not specified on the product(s) label within 12 months (365 days) following a DEVIRINOL 50-DF treatment at rates not exceeding 2 lbs ai/A.

WEEDS CONTROLLED BY 2-4 LB DEVIRINOL 50-DF (1-2 LB. AI/ACRE)

NOTE: DEVIRINOL does not control established weeds.

Barnyardgrass	Junglerice
Carpetweed	Lambsquarters
Common Purslane	Panicum
Crabgrass	Purple Cudweed
Fall Panicum	Redroot Pigweed
Florida Pusley	Smooth Pigweed
Giant Foxtail	Stinkgrass
Goosegrass	Witchgrass
Green Foxtail	Yellow Foxtail
Johnsongrass (seedling)	

WEEDS CONTROLLED BY 4 LB DEVIRINOL 50-DF (2 LB AI/ACRE)

NOTE: DEVIRINOL does not control established weeds.

Galinsoga

24(c) Registrant:



United Phosphorus, Inc.
630 Freedom Business Center
King of Prussia, PA 19406
1-800-438-6071 • www.upi-usa.com

Rev. 12/2/10