



Recommendation

FIFRA SECTION 2(ee) RECOMMENDATION ONLY FOR DISTRIBUTION IN FLORIDA.

**MICROTHIOL® DISPERSS®
EPA Reg. No. 70506-187**

FOR SUPPRESSION OF CITRUS CANKER IN CITRUS

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by USEPA.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read and follow all applicable instructions, restrictions, and precautions on the EPA-approved label. This label and the federally registered label must be in the possession of the user at the time of pesticide application.

CROP	PEST	DIRECTIONS
Citrus	Citrus Canker (Suppression Only)	Use the labeled rate of a fixed copper fungicide product, such as Cuprofix Ultra 40 Disperss (EPA Reg. No. 70506-201) in a tank mix combination with 3.0-6.0 lbs. of Microthiol Disperss. Spray flushes 7 to 14 days after shoots begin to grow. Young fruit may require an additional application. Number and timing of applications will be dependent upon disease pressure. Under heavy pressure, each flush of new growth should be sprayed.

Do not apply within 21 days of an oil spray.
The addition of adjuvants (spreaders, surfactants, etc.) can increase the potential of crop injury and is not recommended.

Always read and follow the most restrictive label instructions.

Rev. 07/21/2015