



Recommendation

FIFRA SECTION 2(ee) RECOMMENDATION FOR DISTRIBUTION AND IN ARIZONA AND CALIFORNIA

METEOR FUNGICIDE EPA Reg. No. 70506-243

FOR SUPPRESSION OF SCLEROTINIA SPP. IN LETTUCE (Head & Leaf Types)

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by USEPA.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read and follow all applicable instructions, restrictions, and precautions on the EPA-approved label. This label and the federally registered label must be in the possession of the user at the time of pesticide application.

DISEASE	RATE/ACRE	INSTRUCTIONS
<i>Sclerotinia</i> spp.	1.5 – 2.0 pts./A	<p>Apply in a vertical band below the seed line immediately before planting.</p> <p>Use a modified fertilizer shank with the main orifice closed and with supplemental orifices evenly spaced approximately 1-2 inches apart in order to distribute the solution evenly in a vertical band from the soil surface to 6 inches below the soil surface.</p> <p>Do not apply deeper than 6 inches. Apply the solution immediately below the seed line in the root zone of the seedlings. Use a minimum of 20 gallons solution per acre. When applying below the seed line, do not reduce the acre rate.</p> <p>One modified shank application can be made before planting. Up to 2 additional bands or broadcast applications can be made to lettuce and the soil surface as described in the label directions of the EPA-approved label.</p>

RESTRICTIONS

Do not make more than one modified shank treatment or more than 3 applications to any lettuce crop.
PreHarvest Interval (PHI) = 14 days.
Do not cultivate after application. If necessary, make an application during or immediately after cultivation.
Do not drench.

Rev. 9/16/2013